



Mercy Hospital of Pittsburgh

Department of Emergency Medicine

CVA Orders

use after Brain Attack Orders if CT negative for bleed for use in DEM **only**


Page 1 of 2

Mandatory

Draw a line through, initial and date all changes.

If generically equivalent medication is not acceptable., then write "brand necessary."

Circles: one and only one. Squares: one, many, or none.



Initial Orders

Continue all previous DEM Brain Attack Orders

Vital Signs

Vital Signs and mini NIH Stroke Scale Q 5 minutes x3

Consider Systemic Thrombolysis

Clearly defined onset of symptoms within the past 3 hours?

Yes Complete **Thrombolysis Checklist**, below

Unknown

No Skip to **CVA Treatment**, bottom of page 2 §

Thrombolysis Checklist: Absolute Criteria

All must be checked to give systemic thrombolytics. IF unable to check any box, thrombolysis is not appropriate: skip to next page.

- Acute ischemic CVA
- Age > 18 years
- No seizure at onset/reason to suspect SAH
- No known bleeding diathesis
- No LMWH or heparin with increased PTT
- No neurosurgery/major head trauma in 3 mo.
- No CNS aneurysm, AVM, bleed or neoplasm
- No major surgery or trauma within 14 days
- No LP within 7 days
- No active GI/GU bleeding within 21 days
- BP ≤ 185/110 now
- Platelets > 100,000
- INR < 1.8
- No hemorrhage on CT read by neuroradiologist

Thrombolysis Checklist: Relative Criteria

If cannot check all, relative contraindication to systemic thrombolysis

- Not rapidly improving neurological status
- Glucose 50-400
- Not pregnant
- No early signs of infarct (edema, sulcal effacement) on CT

Continued next page...

Resident/PA/CRNP:

Attending emergency physician (required):

Print Last Name:

Print Last Name:

Date:

Sign:

Date:

Sign:

Time:

Time: